



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/31, C12N 1/21, 15/31, 15/63, 15/74	A1	(11) International Publication Number: WO 99/31133 (43) International Publication Date: 24 June 1999 (24.06.99)
(21) International Application Number: PCT/US98/26037 (22) International Filing Date: 8 December 1998 (08.12.98) (30) Priority Data: 08/993,825 18 December 1997 (18.12.97) US (71) Applicant: SMITHKLINE BEECHAM CORPORATION [US/US]; One Franklin Plaza, Philadelphia, PA 19103 (US). (72) Inventors: WARREN, Richard, L.; 825 Cathcart Road, Blue Bell, PA 19422 (US). TRAINI, Christopher, M.; 50 Potter Court, Media, PA 19063 (US). (74) Agent: FISH, Andrew, L.; Dechert Price & Rhoads, 4000 Bell Atlantic Tower, 1717 Arch Street, Philadelphia, PA 19103 (US).		(81) Designated States: JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: NOVEL ABC TRANSPORTER (57) Abstract The invention provides Novel ABC transporter polypeptides and polynucleotides encoding Novel ABC transporter polypeptides and methods for producing such polypeptides by recombinant techniques. Also provided are methods for utilizing Novel ABC transporter polypeptides to screen for antibacterial compounds.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

Novel ABC transporter

FIELD OF THE INVENTION

5 This invention relates to newly identified polynucleotides and polypeptides, and their production and uses, as well as their variants, agonists and antagonists, and their uses. In particular, the invention relates to novel polynucleotides and polypeptides of the ABC transporter family, hereinafter referred to as "Novel ABC transporter".

BACKGROUND OF THE INVENTION

10

It is particularly preferred to employ Staphylococcal genes and gene products as targets for the development of antibiotics. The Staphylococci make up a medically important genera of microbes. They are known to produce two types of disease, invasive and toxigenic. Invasive infections are characterized generally by abscess formation effecting both skin surfaces and deep tissues. *Staphylococcus aureus* is the second leading cause of bacteremia in cancer patients. Osteomyelitis, septic arthritis, septic thrombophlebitis and acute bacterial endocarditis are also relatively common. There are at least three clinical conditions resulting from the toxigenic properties of Staphylococci. The manifestation of these diseases result from the actions of exotoxins as opposed to tissue invasion and bacteremia. These conditions include: Staphylococcal food poisoning, scalded skin syndrome and toxic shock syndrome.

20

The frequency of *Staphylococcus aureus* infections has risen dramatically in the past few decades. This has been attributed to the emergence of multiply antibiotic resistant strains and an increasing population of people with weakened immune systems. It is no longer uncommon to isolate *Staphylococcus aureus* strains which are resistant to some or all of the standard antibiotics. This phenomenon has created a demand for both new anti-microbial agents, vaccines, and diagnostic tests for this organism.

25

Clearly, there exists a need for factors, such as the Novel ABC transporter embodiments of the invention, that have a present benefit of being useful to screen compounds for antibiotic activity. Such factors are also useful to determine their role in pathogenesis of infection, dysfunction and disease. There is also a need for identification and characterization of such factors and their

30

antagonists and agonists to find ways to prevent, ameliorate or correct such infection, dysfunction and disease.

Certain of the polypeptides of the invention possess amino acid sequence homology to a known *Archaeoglobus fulgidus* ABC transporter (AE001033) protein. (Higgins et al. 1990. J. Bioenerg. Biomembr 22:571-592; Aguilar_Bryan et al 1995 Science 268:423-426. Rosteck et al 1991 Gene 102:27-32; Accession numbers AE001023, C64488.)

SUMMARY OF THE INVENTION

5 It is an object of the invention to provide polypeptides that have been identified as novel Novel ABC transporter polypeptides by homology between the amino acid sequence set out in Table 1 [SEQ ID NO: 2] and a known amino acid sequence or sequences of other proteins such as *Archaeoglobus fulgidus* ABC transporter (AE001033) protein.

10 It is a further object of the invention to provide polynucleotides that encode Novel ABC transporter polypeptides, particularly polynucleotides that encode the polypeptide herein designated Novel ABC transporter.

In a particularly preferred embodiment of the invention, the polynucleotide comprises a region encoding Novel ABC transporter polypeptides comprising a sequence set out in Table 1 [SEQ ID NO:1] which includes a full length gene, or a variant thereof.

15 In another particularly preferred embodiment of the invention,, there is a novel Novel ABC transporter protein from *Staphylococcus aureus* comprising the amino acid sequence of Table 1 [SEQ ID NO:2], or a variant thereof.

20 As a further aspect of the invention there are provided isolated nucleic acid molecules encoding Novel ABC transporter, particularly *Staphylococcus aureus* Novel ABC transporter, including mRNAs, cDNAs, genomic DNAs. Further embodiments of the invention include biologically, diagnostically, prophylactically, clinically or therapeutically useful variants thereof, and compositions comprising the same.

25 In accordance with another aspect of the invention, there is provided the use of a polynucleotide of the invention for therapeutic or prophylactic purposes, in particular genetic immunization. Among the particularly preferred embodiments of the invention are naturally occurring allelic variants of Novel ABC transporter and polypeptides encoded thereby.

As another aspect of the invention, there are provided novel polypeptides of *Staphylococcus aureus* referred to herein as Novel ABC transporter as well as biologically, diagnostically, prophylactically, clinically or therapeutically useful variants thereof, and compositions comprising the same.

5 Among the particularly preferred embodiments of the invention are variants of Novel ABC transporter polypeptide encoded by naturally occurring alleles of the Novel ABC transporter gene.

In a preferred embodiment of the invention, there are provided methods for producing the aforementioned Novel ABC transporter polypeptides.

10 In accordance with yet another aspect of the invention, there are provided inhibitors to such polypeptides, useful as antibacterial agents, including, for example, antibodies.

In accordance with certain preferred embodiments of the invention, there are provided products, compositions and methods for assessing Novel ABC transporter expression, treating disease, assaying genetic variation, and administering a Novel ABC transporter polypeptide or polynucleotide to an organism to raise an immunological response against a bacteria, especially a
15 *Staphylococcus aureus* bacteria.

In accordance with certain preferred embodiments of this and other aspects of the invention, there are provided polynucleotides that hybridize to Novel ABC transporter polynucleotide sequences, particularly under stringent conditions.

20 In certain preferred embodiments of the invention, there are provided antibodies against Novel ABC transporter polypeptides.

In other embodiments of the invention, there are provided methods for identifying compounds which bind to or otherwise interact with and inhibit or activate an activity of a polypeptide or polynucleotide of the invention comprising: contacting a polypeptide or polynucleotide of the invention with a compound to be screened under conditions to permit binding to or other
25 interaction between the compound and the polypeptide or polynucleotide to assess the binding to or other interaction with the compound, such binding or interaction being associated with a second component capable of providing a detectable signal in response to the binding or interaction of the polypeptide or polynucleotide with the compound; and determining whether the compound binds to or otherwise interacts with and activates or inhibits an activity of the polypeptide or polynucleotide
30 by detecting the presence or absence of a signal generated from the binding or interaction of the compound with the polypeptide or polynucleotide.

In accordance with yet another aspect of the invention, there are provided Novel ABC transporter agonists and antagonists, preferably bacteriostatic or bacteriocidal agonists and antagonists.

In a further aspect of the invention, there are provided compositions comprising a Novel ABC transporter polynucleotide or a Novel ABC transporter polypeptide for administration to a cell or to a multicellular organism.

Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following descriptions and from reading the other parts of the present disclosure.

DESCRIPTION OF THE INVENTION

The invention relates to novel Novel ABC transporter polypeptides and polynucleotides as described in greater detail below. In particular, the invention relates to polypeptides and polynucleotides of a novel Novel ABC transporter of *Staphylococcus aureus*, which is related by amino acid sequence homology to *Archaeoglobus fulgidus* ABC transporter (AE001033) polypeptide. The invention relates especially to Novel ABC transporter having the nucleotide and amino acid sequences set out in Table 1 as SEQ ID NO: 1 and SEQ ID NO: 2, respectively.

20

TABLE 1

Novel ABC transporter Polynucleotide and Polypeptide Sequences

(A) Sequences from *Staphylococcus aureus* Novel ABC transporter polynucleotide sequence [SEQ ID NO:1].

25

5' -CATTATATATTGTTGTCTTAATGTTTTATCAGAAGTTTGTTGTAACG
CTTAATAAATGATGAATATGTGTTAATGGTGTGTTTGATTTTCATGAGATAC
ATTTTGTAACAAATGCTGCCATTTGATCAACTTGCCCAATGACTCTC
TCATCTTATTAAAGTGATACTGTAATGTTCCAATTTTCATCTTTGCGTGTT
TGTTTGATAGGTGTTTCAAAATCACCATCAATTAAGCGTTCGGTCGCTAG
30 CTTTAACTTTTTTACTGGACGAATGATTGAATAGGTTGATGCGATAACTA
ATGAAATCGAGATAAATAACAACATTAACAAGACGGCTAAGAATGTT
CTAAACTCGCTAAATGTTTCTCCAATATCTGGACGCATGAATACTGCTAT
TGAACCGTCTTTTGTCTTAAAGTTAATACCAACAGTATTATCTGTTACAT
TGTCGAAAAATCCAGTTACGAATAATGCGAATGTTTATCTTTAATACCAT

GGTAATCTTTATTATTTAATACGTTTGTAATAGCATTTTGTGAAAAAGTA
TCTTCTCTAAATGGCTCCCCAAAAATGTCTTATGACCCTTTTGATCAACG
GTCATAATTTGGTAATTCATTGGCCCTAAATGTTTGAAATATTGTTGAAT
GTGCGTTGGTTTAACTGATTGTTTCATATTGTCTTGCTTCTTTAAGTGTCT
5 TCATGATTTTCGCGTCATTAGATGCTTTTAAATTATAATGATAGTAAACA
TTTGTTAATACAAAACCTTATTAATGCACTAAATAAAATAACCGTAATGGA
ATAAATCGCAATTCATAGCATAGAGTGTTTTAAACATGATTCTCCACCTTA
TAGCCTTGTCCTCTTACTGTTTCAATTGTAAGTGTGGCATTTAATTTTTT
TAATCTTTGGCGTAGTCGCTTAATATGAACGTCAACTGTTGTTTCATCTC
10 CTTCATAATCATAGCCCCAAATTTTTCTATTATTTGTTACGAGTAAAG
ATTTGCTTAGGACGTGCTGCAAGCATAAATAAATTGAAATTCCTTGTT
CGGTAACGTCATCGTTTATTAGATACTTGAGATTCCAAATAGGATTGGTT
TAGCGTTAAGTTGCCAATAGTCATTTCTGAATTTGAATTGATATTATATC
GACGTAATACCGCACGAATTCTAAAAATAAGTTCCTTAACCTCAAAGGGT
15 TTGGTTACATAATCGTCAGTACCACTTATAAACGCACGCTCTTTGTCACT
AAGTGCATCCCGCGCTGTTAACATAATAACTGGTATATCATAATCATTTT
TTAATGTATTACATAATTGAAAGCCGTCCATACCATCCATCATAATATCT
ACCACTGCAATATCGACACGCTGTTTTCTAAAAGTTTTAATGCAGCTTC
ACCACTTGTTGTGTGTATGCATCAATGTGCTCTGTTTGTAATGGCTAG
20 CTATATAATTTAGGATTCGAGGATCGTCATCGACAACAAGACATTGCACC
ATAGCTATAAACTCCCTTATCTTTTTCATTTATTATACATGTAAATATT
TTTGCGTAAAAAACCAATTGTTTCATATTGAGTTCATATTTCAACCTTATA
CTGACGCTAAAGAAGAAATAGGGAGAAGTGAATCGATATGAAATTAGCGA
TAAAAGAGATTATGTTTACAAATTCGTTATATTTAATCACATTAATC
25 ATTCCTTTTATTAAGTATTATGGGGTTATTTATTAGTGGTTTAACTCAAGG
GCTTGGTAGGGAGAATATTCGTTATTTGAACACTTTGATAATGAATAAT
ATGTTGTTCAAAAAATGAAAGAGCCGCAAAATTGAGAAATCGCAACTTAGT
GATACGCAACAGAATCAATTAAAAAAGTAATTCATCAAGAACCATATAAA
ATGAATATTCAAACATTGAAATTATCTAATAAAGAACAAGATGTGATTAC
30 ATGAATGACGTTAAACAACAACGCCTACAACCTAAAAAAGGTGAATATCC
TAAAAATGCTCATGAGGTCGCTATTAATGATAAGTTAGCTGCAGACAACA
TTAGAGTCGGGGATAGATTACATTTTAAAAATAATTCAACTAGTTATAGA
GTTTCTGGTATTTTAAACGACACAATGTATGCGCATAGTTCCATTGTGCT
ATTGAACGATAACGGATTTAATGCATTGAATAAGGTAAATACGGCATTTT
35 ATCCAGTGAAAAATTTAACACAACAACAACGTGATGAGCTTAATAAAATA
AATGACGTTCAAGTTGTGAGTGAAAAAGATTTAACAGGTAATATTGCGAG
TTATCAAGCAGAGCAAGCACCGTTAAATATGATGATTGTTAGTTTGTTTG
CTATTACAGCAATCGTTCTAAGTGCATTTTTCTATGTTATGACGATTCAA
AAAATATCACAAATTGGCATTTTGAAAGCAATTGGTATTAAGACAAGACA

TTTATTGAGTGCCTTAGTTTTACAAATTTTAACTAACAATAATTGGGG
TAGGTATTGCTGTGATCATCATAGTAGGACTATCATTTATGATGCCGGTA
ACGATGCCTTTTTTACTTAAACAACGCAAAATATTTTATTAATGGTGGGGAT
ATTTATATTAGTAGCGATTTTAGGTGCCTCACTATCATTTATCAAATTAT
5 TTAAGTGGATCCTATCGAAGCAATTGGAGGTGCAGAATAATGGCATTAG
TCGTTAAAGATATCGTCAAAAATTTTCGGAGAAGGTTTGTCTGAAACAAA
GTTTTAAAGGTATTAATTTTGAAGTGAACAAGGGGAATTTGTCATTTT
AAATGGTGCCTCTGGTTCTGGGAAAACAACATTGCTAACGATATTAGGCG
GATTGTTAAGTCAAACGAGTGGTACAGTGCTTTACAATGATGCACCATTG
10 TTTGATAAACAGCATCGTCCTAGTGATTTGCGATTGGAAGATATTGGTTT
TATTTTTCAATCTTCACATTTAGTTCCTTATTTAAAGTGATAGAGCAAT
TGACACTCGTAGGCCAAGAAGCGGGGAATGACCAACAACAAAGTCAACA
AGAGCAATACAACTTTTGAAAAATATTGGGTTAGAAGATCGCTGAATGT
ATATCCGCATCAGTTATCTGGCGGTGAAAAGCAACGTGTTGCGATTATGA
15 GAGCATTTATGAATAATCCGAAAATCATTTTAGCAGATGAGCCACAGCA
AGTTTAGATGCCGATAGAGCAACAAAAGTTGTTGAGATGATACGTCAACA
AATTAAAGAACAACAATGATTGGTATTATGATTACACACGATCGAAGAT
TATTTGAATATGCAGATCGAGTGATTGAATTAGAAGATGGCAAATAACT
GATTAGTGGCTTGTAAGACGCTAAATGTTAATGATTTAAGACATAGTAG
20 TATAAAGTTAGATAACAGAATACGATTTGGGTTTACAAAAAGCGCGTC
ACTATTATATAAAATGATGACGTGCTTTAGTTTATTTATTTTCAATATAA
CAATTGTCTAATCAAGGATTAACGAATATTTAAAGATAGTTTGACGCAAT
ATTAGAAACAACCTATAATAATAGTTTGTGTTGGGATTAATATTATAAA
TAAAAGCGGCGTAAAGACATATAAACCAACTACTTGAACAATATAACGTT
25 AATAACAATCTATACTGATACATTACGCCTAGATAATCTTTAATGAGCAC
ATGTAAGAAAAAGTGATATGGTGTATGACTTCCGACACCATCGATAGATA
AACCTAATTTTTGGGCTAGTCGTAAGGCGCGCAATACATGAACTGACTT
GTTACACAAACAATTTTAACTGCTTCATGATACAAATTGTTGATGATTG
TTTAGAATATAAAAAGTTTGTGTATGTATTTATAGAGTGAGATTCCATTA
30 GTATATCTGTTTTATCAACACCATGTGCAATCAAATAACGTTGCATAGCT
AAAGCTTCAGAAATTGGTTCGTCTGGTCCTTGTCGCCAGATACAATAAT
CTTTGTTGCTGATGCTTGTGTTGATAGATATCAAGTGCACGGTCTAAAC
GCGCTGCAAGCATTGGTGTGACAAATTCGGTAAAAATACCAGCACCTAAC
ACAATTATGATATCAACTTCTTGTGTTGATGATCTATGTCTATATGATAC
35 TGTCCAAACGAGATAACAAATAAAGGTTAGTAACAGGGAAAGACATAATA
TAGCTAACCACATAGACAAACCTTTACAATAGGTGACTGAATCGTACTT
ATAAATAGAAGTGCTGATGTGTAGAGTAC-3'

(B) *Staphylococcus aureus* Novel ABC transporter polypeptide sequence deduced from the polynucleotide sequence in this table [SEQ ID NO:2].

NH₂-

SGSYRSNWRCRIMALVVKDIVKNFGEGLSETKVLKGINFEVEQGEFVILNGASGSGKTTLLTILGGLLS
 5 QTSGLTVLYNDAPLEFDKQHRPSDLRLEDIGFIFQSSHLVPYLKVIEQLTLVGQEAGMTKQSSSTRAIQLL
 KNIGLEDRLNVYPHQLSGGEKQRVAIMRAFMNNPKIILADEPTASLDADRATKVVEMIRQQIKEQQMIG
 IMITHDRRLFEYADRVIELEDGKITD-COOH

Deposited materials

10 A deposit containing a *Staphylococcus aureus* WCUH 29 strain has been deposited with the National Collections of Industrial and Marine Bacteria Ltd. (herein "NCIMB"), 23 St. Machar Drive, Aberdeen AB2 1RY, Scotland on 11 September 1995 and assigned NCIMB Deposit No. 40771, and referred to as *Staphylococcus aureus* WCUH29 on deposit. The *Staphylococcus aureus* strain deposit is referred to herein as "the deposited strain" or as "the DNA of the deposited strain."

15 The deposited strain contains the full length Novel ABC transporter gene. The sequence of the polynucleotides contained in the deposited strain, as well as the amino acid sequence of the polypeptide encoded thereby, are controlling in the event of any conflict with any description of sequences herein.

The deposit of the deposited strain has been made under the terms of the Budapest Treaty
 20 on the International Recognition of the Deposit of Micro-organisms for Purposes of Patent Procedure. The strain will be irrevocably and without restriction or condition released to the public upon the issuance of a patent. The deposited strain is provided merely as convenience to those of skill in the art and is not an admission that a deposit is required for enablement, such as that required under 35 U.S.C. §112.


25 A license may be required to make, use or sell the deposited strain, and compounds derived therefrom, and no such license is hereby granted.

One aspect of the invention there is provided an isolated nucleic acid molecule encoding a mature polypeptide expressible by the *Staphylococcus aureus* WCUH 29 strain contained in the deposited strain. Further provided by the invention are Novel ABC transporter nucleotide sequences
 30 of the DNA in the deposited strain and amino acid sequences encoded thereby. Also provided by the invention are Novel ABC transporter polypeptide sequences isolated from the deposited strain and amino acid sequences derived therefrom.

Polypeptides

The polypeptides of the invention include a polypeptide of Table 1 [SEQ ID NO:2] (in particular the mature polypeptide) as well as polypeptides and fragments, particularly those which have the biological activity of Novel ABC transporter, and also those which have at least 70% identity to a polypeptide of Table 1 [SEQ ID NO:1] or the relevant portion, preferably at least 80% identity to a polypeptide of Table 1 [SEQ ID NO:2] and more preferably at least 90% similarity (more preferably at least 90% identity) to a polypeptide of Table 1 [SEQ ID NO:2] and still more preferably at least 95% similarity (still more preferably at least 95% identity) to a polypeptide of Table 1 [SEQ ID NO:2] and also include portions of such polypeptides with such portion of the polypeptide generally containing at least 30 amino acids and more preferably at least 50 amino acids.

The invention also includes polypeptides of the formula:



wherein, at the amino terminus, X is hydrogen, and at the carboxyl terminus, Y is hydrogen or a metal, R_1 and R_3 are any amino acid residue, m is an integer between 1 and 1000 or zero, n is an integer between 1 and 1000 or zero, and R_2 is an amino acid sequence of the invention, particularly an amino acid sequence selected from Table 1. In the formula above R_2 is oriented so that its amino terminal residue is at the left, bound to R_1 , and its carboxy terminal residue is at the right, bound to R_3 . Any stretch of amino acid residues denoted by either R group, where m and/or n is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer.

A fragment is a variant polypeptide having an amino acid sequence that entirely is the same as part but not all of the amino acid sequence of the aforementioned polypeptides. As with Novel ABC transporter polypeptides fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most preferably as a single continuous region, a single larger polypeptide.

Preferred fragments include, for example, truncation polypeptides having a portion of an amino acid sequence of Table 1 [SEQ ID NO:2], or of variants thereof, such as a continuous series of residues that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus. Degradation forms of the polypeptides of the invention in a host cell, particularly a *Staphylococcus aureus*, are also preferred. Further preferred are fragments characterized by structural or functional attributes such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta

amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions.

Also preferred are biologically active fragments which are those fragments that mediate activities of Novel ABC transporter, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also included are those fragments that are antigenic or immunogenic in an animal, especially in a human. Particularly preferred are fragments comprising receptors or domains of enzymes that confer a function essential for viability of *Staphylococcus aureus* or the ability to initiate, or maintain cause disease in an individual, particularly a human.

Variants that are fragments of the polypeptides of the invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, these variants may be employed as intermediates for producing the full-length polypeptides of the invention.

In addition to the standard single and triple letter representations for amino acids, the term "X" or "Xaa" may also be used in describing certain polypeptides of the invention. "X" and "Xaa" mean that any of the twenty naturally occurring amino acids may appear at such a designated position in the polypeptide sequence.

Polynucleotides

Another aspect of the invention relates to isolated polynucleotides, including the full length gene, that encode the Novel ABC transporter polypeptide having a deduced amino acid sequence of Table 1 [SEQ ID NO:2] and polynucleotides closely related thereto and variants thereof.

Using the information provided herein, such as a polynucleotide sequence set out in Table 1 [SEQ ID NO:1], a polynucleotide of the invention encoding Novel ABC transporter polypeptide may be obtained using standard cloning and screening methods, such as those for cloning and sequencing chromosomal DNA fragments from bacteria using *Staphylococcus aureus* WCUH 29 cells as starting material, followed by obtaining a full length clone. For example, to obtain a polynucleotide sequence of the invention, such as a sequence given in Table 1 [SEQ ID NO:1], typically a library of clones of chromosomal DNA of *Staphylococcus aureus* WCUH 29 in *E.coli* or some other suitable host is probed with a radiolabeled oligonucleotide, preferably a 17-mer or longer, derived from a partial sequence. Clones carrying DNA identical to that of the probe can then be distinguished using stringent conditions. By sequencing the individual clones thus identified with sequencing primers designed from the original sequence it is then possible to extend the sequence in both directions to determine the full gene sequence. Conveniently, such sequencing is performed using denatured double stranded DNA prepared from a plasmid clone. Suitable

techniques are described by Maniatis, T., Fritsch, E.F. and Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989). (see in particular Screening By Hybridization 1.90 and Sequencing Denatured Double-Stranded DNA Templates 13.70). Illustrative of the invention,
5 the polynucleotide set out in Table 1 [SEQ ID NO:1] was discovered in a DNA library derived from *Staphylococcus aureus* WCUH 29.


The DNA sequence set out in Table 1 [SEQ ID NO:1] contains an open reading frame encoding a protein having about the number of amino acid residues set forth in Table 1 [SEQ ID NO:2] with a deduced molecular weight that can be calculated using amino acid residue molecular
10 weight values well known in the art. The polynucleotide of SEQ ID NO: 1, between nucleotide number 2655 and the stop codon which begins at nucleotide number 3354 of SEQ ID NO:1, encodes the polypeptide of SEQ ID NO:2.

Novel ABC transporter of the invention is structurally related to other proteins of the ABC transporter family.

15 The invention provides a polynucleotide sequence identical over its entire length to a coding sequence in Table 1 [SEQ ID NO:1]. Also provided by the invention is the coding sequence for the mature polypeptide or a fragment thereof, by itself as well as the coding sequence for the mature polypeptide or a fragment in reading frame with other coding sequence, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence. The polynucleotide may
20 also contain non-coding sequences, including for example, but not limited to non-coding 5' and 3' sequences, such as the transcribed, non-translated sequences, termination signals, ribosome binding sites, sequences that stabilize mRNA, introns, polyadenylation signals, and additional coding sequence which encode additional amino acids. For example, a marker sequence that facilitates purification of the fused polypeptide can be encoded. In certain embodiments of the invention, the
25 marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, *Proc. Natl. Acad. Sci., USA* 86: 821-824 (1989), or an HA tag (Wilson *et al.*, *Cell* 37: 767 (1984). Polynucleotides of the invention also include, but are not limited to, polynucleotides comprising a structural gene and its naturally associated sequences that control gene expression.

30 A preferred embodiment of the invention is a polynucleotide of comprising nucleotide 2655 to the nucleotide immediately upstream of or including nucleotide 3354 set forth in SEQ ID NO:1 of Table 1, both of which encode the Novel ABC transporter polypeptide.

The invention also includes polynucleotides of the formula:



wherein, at the 5' end of the molecule, X is hydrogen or together with Y defines a covalent bond, and at the 3' end of the molecule, Y is hydrogen or a metal or together with X defines the covalent bond, each occurrence of R_1 and R_3 is independently any nucleic acid residue, m is an integer between 1 and 3000 or zero, n is an integer between 1 and 3000 or zero, and R_2 is a nucleic acid sequence of the invention, particularly a nucleic acid sequence selected from Table 1. In the polynucleotide formula above R_2 is oriented so that its 5' end residue is at the left, bound to R_1 , and its 3' end residue is at the right, bound to R_3 . Any stretch of nucleic acid residues denoted by either R group, where m and/or n is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer. Where, in a preferred embodiment, X and Y together define a covalent bond, the polynucleotide of the above formula is a closed, circular polynucleotide, which can be a double-stranded polynucleotide wherein the formula shows a first strand to which the second strand is complementary. In another preferred embodiment m and/or n is an integer between 1 and 1000.

It is most preferred that the polynucleotides of the inventions are derived from *Staphylococcus aureus*, however, they may preferably be obtained from organisms of the same taxonomic genus. They may also be obtained, for example, from organisms of the same taxonomic family or order.

The term "polynucleotide encoding a polypeptide" as used herein encompasses polynucleotides that include a sequence encoding a polypeptide of the invention, particularly a bacterial polypeptide and more particularly a polypeptide of the *Staphylococcus aureus* Novel ABC transporter having an amino acid sequence set out in Table 1 [SEQ ID NO:2]. The term also encompasses polynucleotides that include a single continuous region or discontinuous regions encoding the polypeptide (for example, interrupted by integrated phage or an insertion sequence or editing) together with additional regions, that also may contain coding and/or non-coding sequences.

The invention further relates to variants of the polynucleotides described herein that encode for variants of the polypeptide having a deduced amino acid sequence of Table 1 [SEQ ID NO:2]. Variants that are fragments of the polynucleotides of the invention may be used to synthesize full-length polynucleotides of the invention.

Further particularly preferred embodiments are polynucleotides encoding Novel ABC transporter variants, that have the amino acid sequence of Novel ABC transporter polypeptide of

Table 1 [SEQ ID NO:2] in which several, a few, 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues are substituted, deleted or added, in any combination. Especially preferred among these are silent substitutions, additions and deletions, that do not alter the properties and activities of Novel ABC transporter.

5 Further preferred embodiments of the invention are polynucleotides that are at least 70% identical over their entire length to a polynucleotide encoding Novel ABC transporter polypeptide having an amino acid sequence set out in Table 1 [SEQ ID NO:2], and polynucleotides that are complementary to such polynucleotides. Alternatively, most highly preferred are polynucleotides that comprise a region that is at least 80% identical over its entire length to a polynucleotide
10 encoding Novel ABC transporter polypeptide and polynucleotides complementary thereto. In this regard, polynucleotides at least 90% identical over their entire length to the same are particularly preferred, and among these particularly preferred polynucleotides, those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred among those with at least 95%, and among these those with at least 98% and at least 99% are particularly highly
15 preferred, with at least 99% being the more preferred.

Preferred embodiments are polynucleotides that encode polypeptides that retain substantially the same biological function or activity as the mature polypeptide encoded by a DNA of Table 1 [SEQ ID NO:1].

The invention further relates to polynucleotides that hybridize to the herein above-described
20 sequences. In this regard, the invention especially relates to polynucleotides that hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, the terms "stringent conditions" and "stringent hybridization conditions" mean hybridization will occur only if there is at least 95% and preferably at least 97% identity between the sequences. An example of stringent hybridization conditions is overnight incubation at 42°C in a solution comprising:
25 50% formamide, 5x SSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml denatured, sheared salmon sperm DNA, followed by washing the hybridization support in 0.1x SSC at about 65°C. Hybridization and wash conditions are well known and exemplified in Sambrook, *et al.*, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, N.Y.,
30 (1989), particularly Chapter 11 therein.

The invention also provides a polynucleotide consisting essentially of a polynucleotide sequence obtainable by screening an appropriate library containing the complete gene for a

polynucleotide sequence set forth in SEQ ID NO:1 under stringent hybridization conditions with a probe having the sequence of said polynucleotide sequence set forth in SEQ ID NO:1 or a fragment thereof; and isolating said DNA sequence. Fragments useful for obtaining such a polynucleotide include, for example, probes and primers described elsewhere herein.

5 As discussed additionally herein regarding polynucleotide assays of the invention, for instance, polynucleotides of the invention as discussed above, may be used as a hybridization probe for RNA, cDNA and genomic DNA to isolate full-length cDNAs and genomic clones encoding Novel ABC transporter and to isolate cDNA and genomic clones of other genes that have a high sequence similarity to the Novel ABC transporter gene. Such probes generally will comprise at least
10 15 bases. Preferably, such probes will have at least 30 bases and may have at least 50 bases. Particularly preferred probes will have at least 30 bases and will have 50 bases or less.

 For example, the coding region of the Novel ABC transporter gene may be isolated by screening using a DNA sequence provided in Table 1 [SEQ ID NO: 1] to synthesize an oligonucleotide probe. A labeled oligonucleotide having a sequence complementary to that of a gene
15 of the invention is then used to screen a library of cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

 The polynucleotides and polypeptides of the invention may be employed, for example, as research reagents and materials for discovery of treatments of and diagnostics for disease, particularly human disease, as further discussed herein relating to polynucleotide assays.

20 Polynucleotides of the invention that are oligonucleotides derived from the sequences of Table 1 [SEQ ID NOS:1 or 2] may be used in the processes herein as described, but preferably for PCR, to determine whether or not the polynucleotides identified herein in whole or in part are transcribed in bacteria in infected tissue. It is recognized that such sequences will also have utility in diagnosis of the stage of infection and type of infection the pathogen has attained.

25 The invention also provides polynucleotides that may encode a polypeptide that is the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature polypeptide (when the mature form has more than one polypeptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, may allow protein transport, may lengthen or shorten protein half-life or may facilitate manipulation of a protein
30 for assay or production, among other things. As generally is the case *in vivo*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

A precursor protein, having the mature form of the polypeptide fused to one or more prosequences may be an inactive form of the polypeptide. When prosequences are removed such inactive precursors generally are activated. Some or all of the prosequences may be removed before activation. Generally, such precursors are called proproteins.

5 In addition to the standard A, G, C, T/U representations for nucleic acid bases, the term "N" may also be used in describing certain polynucleotides of the invention. "N" means that any of the four DNA or RNA bases may appear at such a designated position in the DNA or RNA sequence, except it is preferred that N is not a base that when taken in combination with adjacent nucleotide positions, when read in the correct reading frame, would have the effect of generating a premature termination codon in such reading frame.

10 In sum, a polynucleotide of the invention may encode a mature protein, a mature protein plus a leader sequence (which may be referred to as a preprotein), a precursor of a mature protein having one or more prosequences that are not the leader sequences of a preprotein, or a preproprotein, which is a precursor to a proprotein, having a leader sequence and one or more prosequences, which generally are removed during processing steps that produce active and mature forms of the polypeptide.

Vectors, host cells, expression

The invention also relates to vectors that comprise a polynucleotide or polynucleotides of the invention, host cells that are genetically engineered with vectors of the invention and the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the invention.

20 For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof or polynucleotides of the invention. Introduction of a polynucleotide into the host cell can be effected by methods described in many standard laboratory manuals, such as Davis et al., *BASIC METHODS IN MOLECULAR BIOLOGY*, (1986) and Sambrook et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989), such as, calcium phosphate transfection, DEAE-dextran mediated transfection, transfection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction and infection.

30 Representative examples of appropriate hosts include bacterial cells, such as streptococci, staphylococci, enterococci *E. coli*, streptomyces and *Bacillus subtilis* cells; fungal cells, such as

yeast cells and *Aspergillus* cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, 293 and Bowes melanoma cells; and plant cells.

5 A great variety of expression systems can be used to produce the polypeptides of the invention. Such vectors include, among others, chromosomal, episomal and virus-derived vectors, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those
10 derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression system constructs may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides and/or to express a polypeptide in a host may be used for expression in this regard. The appropriate DNA sequence may be inserted into the expression system by any of a variety of
15 well-known and routine techniques, such as, for example, those set forth in Sambrook *et al.*, *MOLECULAR CLONING, A LABORATORY MANUAL*, (*supra*).

For secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the expressed polypeptide. These signals may be endogenous to the polypeptide or
20 they may be heterologous signals.

Polypeptides of the invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography, and lectin
25 chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding protein may be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

Diagnostic Assays

This invention is also related to the use of the Novel ABC transporter polynucleotides of the
30 invention for use as diagnostic reagents. Detection of Novel ABC transporter in a eukaryote, particularly a mammal, and especially a human, will provide a diagnostic method for diagnosis of a disease. Eukaryotes (herein also "individual(s)"), particularly mammals, and especially humans,

particularly those infected or suspected to be infected with an organism comprising the Novel ABC transporter gene may be detected at the nucleic acid level by a variety of techniques.

Nucleic acids for diagnosis may be obtained from an infected individual's cells and tissues, such as bone, blood, muscle, cartilage, and skin. Genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification technique prior to analysis. RNA, cDNA and genomic DNA may also be used in the same ways. Using amplification, characterization of the species and strain of prokaryote present in an individual, may be made by an analysis of the genotype of the prokaryote gene. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the genotype of a reference sequence. Point mutations can be identified by hybridizing amplified DNA to labeled Novel ABC transporter polynucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in the electrophoretic mobility of the DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing. See, *e.g.*, Myers et al., *Science*, 230: 1242 (1985). Sequence changes at specific locations also may be revealed by nuclease protection assays, such as RNase and S1 protection or a chemical cleavage method. See, *e.g.*, Cotton et al., *Proc. Natl. Acad. Sci., USA*, 85: 4397-4401 (1985).


Cells carrying mutations or polymorphisms in the gene of the invention may also be detected at the DNA level by a variety of techniques, to allow for serotyping, for example. For example, RT-PCR can be used to detect mutations. It is particularly preferred to use RT-PCR in conjunction with automated detection systems, such as, for example, GeneScan. RNA, cDNA or genomic DNA may also be used for the same purpose, PCR or RT-PCR. As an example, PCR primers complementary to a nucleic acid encoding Novel ABC transporter can be used to identify and analyze mutations. Examples of representative primers are shown below in Table 2.

Table 2

Primers for amplification of Novel ABC transporter polynucleotides

<u>SEQ ID NO</u>	<u>PRIMER SEQUENCE</u>
3	5'-TGCCTCTGGTTCTGGGAAAA-3'
4	5'-ACTAAATGTGAAGATTGAAAAATAAACCAATA-3'

The invention also includes primers of the formula:



wherein, at the 5' end of the molecule, X is hydrogen, and at the 3' end of the molecule, Y is hydrogen or a metal, R_1 and R_3 is any nucleic acid residue, m is an integer between 1 and 20 or zero, n is an integer between 1 and 20 or zero, and R_2 is a primer sequence of the invention, particularly a primer sequence selected from Table 2. In the polynucleotide formula above R_2 is oriented so that its 5' end residue is at the left, bound to R_1 , and its 3' end residue is at the right, bound to R_3 . Any stretch of nucleic acid residues denoted by either R group, where m and/or n is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer being complementary to a region of a polynucleotide of Table 1. In a preferred embodiment m and/or n is an integer between 1 and 10.

The invention further provides these primers with 1, 2, 3 or 4 nucleotides removed from the 5' and/or the 3' end. These primers may be used for, among other things, amplifying Novel ABC transporter DNA isolated from a sample derived from an individual. The primers may be used to amplify the gene isolated from an infected individual such that the gene may then be subject to various techniques for elucidation of the DNA sequence. In this way, mutations in the DNA sequence may be detected and used to diagnose infection and to serotype and/or classify the infectious agent.

The invention further provides a process for diagnosing, disease, preferably bacterial infections, more preferably infections by *Staphylococcus aureus*, comprising determining from a sample derived from an individual a increased level of expression of polynucleotide having a sequence of Table 1 [SEQ ID NO: 1]. Increased or decreased expression of Novel ABC transporter polynucleotide can be measured using any one of the methods well known in the art for the quantation of polynucleotides, such as, for example, amplification, PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods.

In addition, a diagnostic assay in accordance with the invention for detecting over-expression of Novel ABC transporter protein compared to normal control tissue samples may be used to detect the presence of an infection, for example. Assay techniques that can be used to determine levels of a Novel ABC transporter protein, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

Antibodies

The polypeptides of the invention or variants thereof, or cells expressing them can be used as an immunogen to produce antibodies immunospecific for such polypeptides. "Antibodies" as used herein includes monoclonal and polyclonal antibodies, chimeric, single chain, simianized antibodies and humanized antibodies, as well as Fab fragments, including the products of an Fab immunoglobulin expression library.

Antibodies generated against the polypeptides of the invention can be obtained by administering the polypeptides or epitope-bearing fragments, analogues or cells to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique known in the art that provides antibodies produced by continuous cell line cultures can be used. Examples include various techniques, such as those in Kohler, G. and Milstein, C., *Nature* 256: 495-497 (1975); Kozbor *et al.*, *Immunology Today* 4: 72 (1983); Cole *et al.*, pg. 77-96 in *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc. (1985).

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms such as other mammals, may be used to express humanized antibodies.

Alternatively phage display technology may be utilized to select antibody genes with binding activities towards the polypeptide either from repertoires of PCR amplified v-genes of lymphocytes from humans screened for possessing anti-Noel ABC transporter or from naive libraries (McCafferty, J. *et al.*, (1990), *Nature* 348, 552-554; Marks, J. *et al.*, (1992) *Biotechnology* 10, 779-783). The affinity of these antibodies can also be improved by chain shuffling (Clackson, T. *et al.*, (1991) *Nature* 352, 624-628).

If two antigen binding domains are present each domain may be directed against a different epitope - termed 'bispecific' antibodies.

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptides to purify the polypeptides by affinity chromatography.

Thus, among others, antibodies against Noel ABC transporter- polypeptide may be employed to treat infections, particularly bacterial infections.

Polypeptide variants include antigenically, epitopically or immunologically equivalent variants that form a particular aspect of this invention. The term "antigenically equivalent derivative" as used herein encompasses a polypeptide or its equivalent which will be specifically recognized by certain antibodies which, when raised to the protein or polypeptide according to the invention, interfere with the immediate physical interaction between pathogen

and mammalian host. The term "immunologically equivalent derivative" as used herein encompasses a peptide or its equivalent which when used in a suitable formulation to raise antibodies in a vertebrate, the antibodies act to interfere with the immediate physical interaction between pathogen and mammalian host.

5 The polypeptide, such as an antigenically or immunologically equivalent derivative or a fusion protein thereof is used as an antigen to immunize a mouse or other animal such as a rat or chicken. The fusion protein may provide stability to the polypeptide. The antigen may be associated, for example by conjugation, with an immunogenic carrier protein for example bovine serum albumin (BSA) or keyhole limpet haemocyanin (KLH). Alternatively a multiple
10 antigenic peptide comprising multiple copies of the protein or polypeptide, or an antigenically or immunologically equivalent polypeptide thereof may be sufficiently antigenic to improve immunogenicity so as to obviate the use of a carrier.

Preferably, the antibody or variant thereof is modified to make it less immunogenic in the individual. For example, if the individual is human the antibody may most preferably be
15 "humanized"; where the complementarity determining region(s) of the hybridoma-derived antibody has been transplanted into a human monoclonal antibody, for example as described in Jones, P. et al. (1986), *Nature* 321, 522-525 or Tempest et al., (1991) *Biotechnology* 9, 266-273.

The use of a polynucleotide of the invention in genetic immunization will preferably
20 employ a suitable delivery method such as direct injection of plasmid DNA into muscles (Wolff et al., *Hum Mol Genet* 1992, 1:363, Manthorpe et al., *Hum. Gene Ther.* 1993:4, 419), delivery of DNA complexed with specific protein carriers (Wu et al., *J Biol Chem.* 1989: 264,16985), coprecipitation of DNA with calcium phosphate (Benvenisty & Reshef, *PNAS USA*, 1986:83,9551), encapsulation of DNA in various forms of liposomes (Kaneda et al., *Science*
25 1989:243,375), particle bombardment (Tang et al., *Nature* 1992, 356:152, Eisenbraun et al., *DNA Cell Biol* 1993, 12:791) and *in vivo* infection using cloned retroviral vectors (Seeger et al., *PNAS USA* 1984:81,5849).

Antagonists and agonists - assays and molecules

Polypeptides of the invention may also be used to assess the binding of small molecule
30 substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be

structural or functional mimetics. See, e.g., Coligan *et al.*, *Current Protocols in Immunology* 1(2): Chapter 5 (1991).

The invention also provides a method of screening compounds to identify those which enhance (agonist) or block (antagonist) the action of Novel ABC transporter polypeptides or polynucleotides, particularly those compounds that are bacteriostatic and/or bacteriocidal. The method of screening may involve high-throughput techniques. For example, to screen for agonists or antagonists, a synthetic reaction mix, a cellular compartment, such as a membrane, cell envelope or cell wall, or a preparation of any thereof, comprising Novel ABC transporter polypeptide and a labeled substrate or ligand of such polypeptide is incubated in the absence or the presence of a candidate molecule that may be a Novel ABC transporter agonist or antagonist. The ability of the candidate molecule to agonize or antagonize the Novel ABC transporter polypeptide is reflected in decreased binding of the labeled ligand or decreased production of product from such substrate. Molecules that bind gratuitously, *i.e.*, without inducing the effects of Novel ABC transporter polypeptide are most likely to be good antagonists. Molecules that bind well and increase the rate of product production from substrate are agonists. Detection of the rate or level of production of product from substrate may be enhanced by using a reporter system. Reporter systems that may be useful in this regard include but are not limited to colorimetric labeled substrate converted into product, a reporter gene that is responsive to changes in Novel ABC transporter polynucleotide or polypeptide activity, and binding assays known in the art.

Another example of an assay for Novel ABC transporter antagonists is a competitive assay that combines Novel ABC transporter and a potential antagonist with Novel ABC transporter-binding molecules, recombinant Novel ABC transporter binding molecules, natural substrates or ligands, or substrate or ligand mimetics, under appropriate conditions for a competitive inhibition assay. Novel ABC transporter can be labeled, such as by radioactivity or a colorimetric compound, such that the number of Novel ABC transporter molecules bound to a binding molecule or converted to product can be determined accurately to assess the effectiveness of the potential antagonist.

Potential antagonists include small organic molecules, peptides, polypeptides and antibodies that bind to a polynucleotide or polypeptide of the invention and thereby inhibit or extinguish its activity. Potential antagonists also may be small organic molecules, a peptide, a polypeptide such as a closely related protein or antibody that binds the same sites on a binding molecule, such as a binding molecule, without inducing Novel ABC transporter-induced activities, thereby preventing the action of Novel ABC transporter by excluding Novel ABC transporter from binding.

Potential antagonists include a small molecule that binds to and occupies the binding site of the polypeptide thereby preventing binding to cellular binding molecules, such that normal biological activity is prevented. Examples of small molecules include but are not limited to small organic molecules, peptides or peptide-like molecules. Other potential antagonists include antisense molecules (see Okano, *J. Neurochem.* 56: 560 (1991); *OLIGODEOXYNUCLEOTIDES AS ANTISENSE INHIBITORS OF GENE EXPRESSION*, CRC Press, Boca Raton, FL (1988), for a description of these molecules). Preferred potential antagonists include compounds related to and variants of Novel ABC transporter.

Each of the DNA sequences provided herein may be used in the discovery and development of antibacterial compounds. The encoded protein, upon expression, can be used as a target for the screening of antibacterial drugs. Additionally, the DNA sequences encoding the amino terminal regions of the encoded protein or Shine-Delgarno or other translation facilitating sequences of the respective mRNA can be used to construct antisense sequences to control the expression of the coding sequence of interest.

The invention also provides the use of the polypeptide, polynucleotide or inhibitor of the invention to interfere with the initial physical interaction between a pathogen and mammalian host responsible for sequelae of infection. In particular the molecules of the invention may be used: in the prevention of adhesion of bacteria, in particular gram positive bacteria, to mammalian extracellular matrix proteins on in-dwelling devices or to extracellular matrix proteins in wounds; to block Novel ABC transporter protein-mediated mammalian cell invasion by, for example, initiating phosphorylation of mammalian tyrosine kinases (Rosenshine *et al.*, *Infect. Immun.* 60:2211 (1992); to block bacterial adhesion between mammalian extracellular matrix proteins and bacterial Novel ABC transporter proteins that mediate tissue damage and; to block the normal progression of pathogenesis in infections initiated other than by the implantation of in-dwelling devices or by other surgical techniques.

The antagonists and agonists of the invention may be employed, for instance, to inhibit and treat diseases.

Helicobacter pylori (herein *H. pylori*) bacteria infect the stomachs of over one-third of the world's population causing stomach cancer, ulcers, and gastritis (International Agency for Research on Cancer (1994) Schistosomes, Liver Flukes and *Helicobacter Pylori* (International Agency for Research on Cancer, Lyon, France; <http://www.uicc.ch/ecp/ecp2904.htm>). Moreover, the international Agency for Research on Cancer recently recognized a cause-and-

effect relationship between *H. pylori* and gastric adenocarcinoma, classifying the bacterium as a Group I (definite) carcinogen. Preferred antimicrobial compounds of the invention (agonists and antagonists of Novel ABC transporter) found using screens provided by the invention, particularly broad-spectrum antibiotics, should be useful in the treatment of *H. pylori* infection.

5 Such treatment should decrease the advent of *H. pylori*-induced cancers, such as gastrointestinal carcinoma. Such treatment should also cure gastric ulcers and gastritis.

Vaccines

Another aspect of the invention relates to a method for inducing an immunological response in an individual, particularly a mammal which comprises inoculating the individual

10 with Novel ABC transporter, or a fragment or variant thereof, adequate to produce antibody and/ or T cell immune response to protect said individual from infection, particularly bacterial infection and most particularly *Staphylococcus aureus* infection. Also provided are methods whereby such immunological response slows bacterial replication. Yet another aspect of the invention relates to a method of inducing immunological response in an individual which

15 comprises delivering to such individual a nucleic acid vector to direct expression of Novel ABC transporter, or a fragment or a variant thereof, for expressing Novel ABC transporter, or a fragment or a variant thereof *in vivo* in order to induce an immunological response, such as, to produce antibody and/ or T cell immune response, including, for example, cytokine-producing T cells or cytotoxic T cells, to protect said individual from disease, whether that disease is

20 already established within the individual or not. One way of administering the gene is by accelerating it into the desired cells as a coating on particles or otherwise. Such nucleic acid vector may comprise DNA, RNA, a modified nucleic acid, or a DNA/RNA hybrid.

A further aspect of the invention relates to an immunological composition which, when introduced into an individual capable or having induced within it an immunological response,

25 induces an immunological response in such individual to a Novel ABC transporter or protein coded therefrom, wherein the composition comprises a recombinant Novel ABC transporter or protein coded therefrom comprising DNA which codes for and expresses an antigen of said Novel ABC transporter or protein coded therefrom. The immunological response may be used therapeutically or prophylactically and may take the form of antibody immunity or cellular

30 immunity such as that arising from CTL or CD4+ T cells.

A Novel ABC transporter polypeptide or a fragment thereof may be fused with co-protein which may not by itself produce antibodies, but is capable of stabilizing the first protein

and producing a fused protein which will have immunogenic and protective properties. Thus fused recombinant protein, preferably further comprises an antigenic co-protein, such as lipoprotein D from *Hemophilus influenzae*, Glutathione-S-transferase (GST) or beta-galactosidase, relatively large co-proteins which solubilize the protein and facilitate production and purification thereof. Moreover, the co-protein may act as an adjuvant in the sense of providing a generalized stimulation of the immune system. The co-protein may be attached to either the amino or carboxy terminus of the first protein.

Provided by this invention are compositions, particularly vaccine compositions, and methods comprising the polypeptides or polynucleotides of the invention and immunostimulatory DNA sequences, such as those described in Sato, Y. *et al.* Science 273: 352 (1996).

Also, provided by this invention are methods using the described polynucleotide or particular fragments thereof which have been shown to encode non-variable regions of bacterial cell surface proteins in DNA constructs used in such genetic immunization experiments in animal models of infection with *Staphylococcus aureus* will be particularly useful for identifying protein epitopes able to provoke a prophylactic or therapeutic immune response. It is believed that this approach will allow for the subsequent preparation of monoclonal antibodies of particular value from the requisite organ of the animal successfully resisting or clearing infection for the development of prophylactic agents or therapeutic treatments of bacterial infection, particularly *Staphylococcus aureus* infection, in mammals, particularly humans.

The polypeptide may be used as an antigen for vaccination of a host to produce specific antibodies which protect against invasion of bacteria, for example by blocking adherence of bacteria to damaged tissue. Examples of tissue damage include wounds in skin or connective tissue caused, e.g., by mechanical, chemical or thermal damage or by implantation of indwelling devices, or wounds in the mucous membranes, such as the mouth, mammary glands, urethra or vagina.

The invention also includes a vaccine formulation which comprises an immunogenic recombinant protein of the invention together with a suitable carrier. Since the protein may be broken down in the stomach, it is preferably administered parenterally, including, for example, administration that is subcutaneous, intramuscular, intravenous, or intradermal. Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection

solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the bodily fluid, preferably the blood, of the individual; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

While the invention has been described with reference to certain Novel ABC transporter protein, it is to be understood that this covers fragments of the naturally occurring protein and similar proteins with additions, deletions or substitutions which do not substantially affect the immunogenic properties of the recombinant protein.

Compositions, kits and administration

The invention also relates to compositions comprising the polynucleotide or the polypeptides discussed above or their agonists or antagonists. The polypeptides of the invention may be employed in combination with a non-sterile or sterile carrier or carriers for use with cells, tissues or organisms, such as a pharmaceutical carrier suitable for administration to a subject. Such compositions comprise, for instance, a media additive or a therapeutically effective amount of a polypeptide of the invention and a pharmaceutically acceptable carrier or excipient. Such carriers may include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol and combinations thereof. The formulation should suit the mode of administration. The invention further relates to diagnostic and pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

Polypeptides and other compounds of the invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

The pharmaceutical compositions may be administered in any effective, convenient manner including, for instance, administration by topical, oral, anal, vaginal, intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal or intradermal routes among others.

In therapy or as a prophylactic, the active agent may be administered to an individual as an injectable composition, for example as a sterile aqueous dispersion, preferably isotonic.

Alternatively the composition may be formulated for topical application

for example in the form of ointments, creams, lotions, eye ointments, eye drops, ear drops, mouthwash, impregnated dressings and sutures and aerosols, and may contain appropriate conventional additives, including, for example, preservatives, solvents to assist drug penetration, and emollients in ointments and creams. Such topical formulations may also
5 contain compatible conventional carriers, for example cream or ointment bases, and ethanol or oleyl alcohol for lotions. Such carriers may constitute from about 1% to about 98% by weight of the formulation; more usually they will constitute up to about 80% by weight of the formulation.

For administration to mammals, and particularly humans, it is expected that the daily
10 dosage level of the active agent will be from 0.01 mg/kg to 10 mg/kg, typically around 1 mg/kg. The physician in any event will determine the actual dosage which will be most suitable for an individual and will vary with the age, weight and response of the particular individual. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of
15 this invention.

In-dwelling devices include surgical implants, prosthetic devices and catheters, i.e., devices that are introduced to the body of an individual and remain in position for an extended time. Such devices include, for example, artificial joints, heart valves, pacemakers, vascular grafts, vascular catheters, cerebrospinal fluid shunts, urinary catheters, continuous ambulatory
20 peritoneal dialysis (CAPD) catheters.

The composition of the invention may be administered by injection to achieve a systemic effect against relevant bacteria shortly before insertion of an in-dwelling device. Treatment may be continued after surgery during the in-body time of the device. In addition, the composition could also be used to broaden perioperative cover for any surgical technique to
25 prevent bacterial wound infections, especially *Staphylococcus aureus* wound infections.

Many orthopaedic surgeons consider that humans with prosthetic joints should be considered for antibiotic prophylaxis before dental treatment that could produce a bacteremia. Late deep infection is a serious complication sometimes leading to loss of the prosthetic joint and is accompanied by significant morbidity and mortality. It may therefore be possible to
30 extend the use of the active agent as a replacement for prophylactic antibiotics in this situation.

In addition to the therapy described above, the compositions of this invention may be used generally as a wound treatment agent to prevent adhesion of bacteria to matrix proteins

exposed in wound tissue and for prophylactic use in dental treatment as an alternative to, or in conjunction with, antibiotic prophylaxis.

Alternatively, the composition of the invention may be used to bathe an indwelling device immediately before insertion. The active agent will preferably be present at a concentration of 1 µg/ml to 10mg/ml for bathing of wounds or indwelling devices.

A vaccine composition is conveniently in injectable form. Conventional adjuvants may be employed to enhance the immune response. A suitable unit dose for vaccination is 0.5-5 microgram/kg of antigen, and such dose is preferably administered 1-3 times and with an interval of 1-3 weeks. With the indicated dose range, no adverse toxicological effects will be observed with the compounds of the invention which would preclude their administration to suitable individuals.

Each reference disclosed herein is incorporated by reference herein in its entirety. Any patent application to which this application claims priority is also incorporated by reference herein in its entirety.

15 GLOSSARY

The following definitions are provided to facilitate understanding of certain terms used frequently herein.

"Disease(s)" means and disease caused by or related to infection by a bacteria, including disease, such as, infections of the upper respiratory tract (e.g., otitis media, bacterial tracheitis, acute epiglottitis, thyroiditis), lower respiratory (e.g., empyema, lung abscess), cardiac (e.g., infective endocarditis), gastrointestinal (e.g., secretory diarrhoea, splenic abscess, retroperitoneal abscess), CNS (e.g., cerebral abscess), eye (e.g., blepharitis, conjunctivitis, keratitis, endophthalmitis, preseptal and orbital cellulitis, dacryocystitis), kidney and urinary tract (e.g., epididymitis, intrarenal and perinephric abscess, toxic shock syndrome), skin (e.g., impetigo, folliculitis, cutaneous abscesses, cellulitis, wound infection, bacterial myositis) bone and joint (e.g., septic arthritis, osteomyelitis).

"Host cell" is a cell which has been transformed or transfected, or is capable of transformation or transfection by an exogenous polynucleotide sequence.

"Identity," as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences.

"Identity" and "similarity" can be readily calculated by known methods, including but not limited to those described in (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data*, Part I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., *SIAM J. Applied Math.*, 48: 1073 (1988). Preferred methods to determine identity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are codified in publicly available computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, the GCG program package (Devereux, J., et al., *Nucleic Acids Research* 12(1): 387 (1984)), BLASTP, BLASTN, and FASTA (Atschul, S.F. et al., *J. Molec. Biol.* 215: 403-410 (1990)). The BLAST X program is publicly available from NCBI and other sources (*BLAST Manual*, Altschul, S., et al., NCBI NLM NIH Bethesda, MD 20894; Altschul, S., et al., *J. Mol. Biol.* 215: 403-410 (1990)). As an illustration, by a polynucleotide having a nucleotide sequence having at least, for example, 95% "identity" to a reference nucleotide sequence of SEQ ID NO: 1 it is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence of SEQ ID NO: 1. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence. Analogously, by a polypeptide having an amino acid sequence having at least, for example, 95% identity to a reference amino acid sequence of SEQ ID NO:2 is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids

of the reference amino acid of SEQ ID NO: 2. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

"Isolated" means altered "by the hand of man" from its natural state, *i.e.*, if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living organism is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein. Moreover, a polynucleotide or polypeptide that is introduced into an organism by transformation, genetic manipulation or by any other recombinant method is "isolated" even if it is still present in said organism, which organism may be living or non-living.

"Polynucleotide(s)" generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotide(s)" include, without limitation, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions or single-, double- and triple-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded, or triple-stranded regions, or a mixture of single- and double-stranded regions. In addition, "polynucleotide" as used herein refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The strands in such regions may be from the same molecule or from different molecules. The regions may include all of one or more of the molecules, but more typically involve only a region of some of the molecules. One of the molecules of a triple-helical region often is an oligonucleotide. As used herein, the term "polynucleotide(s)" also includes DNAs or RNAs as described above that contain one or more modified bases. Thus, DNAs or RNAs with backbones modified for stability or for other reasons are "polynucleotide(s)" as that term is intended herein. Moreover, DNAs or RNAs comprising unusual bases, such as inosine, or modified bases, such as tritylated bases, to name just

two examples, are polynucleotides as the term is used herein. It will be appreciated that a great variety of modifications have been made to DNA and RNA that serve many useful purposes known to those of skill in the art. The term "polynucleotide(s)" as it is employed herein embraces such chemically, enzymatically or metabolically modified forms of polynucleotides, as well as the chemical forms of DNA and RNA characteristic of viruses and cells, including, for example, simple and complex cells. "Polynucleotide(s)" also embraces short polynucleotides often referred to as oligonucleotide(s).

"Polypeptide(s)" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds. "Polypeptide(s)" refers to both short chains, commonly referred to as peptides, oligopeptides and oligomers and to longer chains generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene encoded amino acids. "Polypeptide(s)" include those modified either by natural processes, such as processing and other post-translational modifications, but also by chemical modification techniques. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature, and they are well known to those of skill in the art. It will be appreciated that the same type of modification may be present in the same or varying degree at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains, and the amino or carboxyl termini. Modifications include, for example, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins, such as arginylation, and ubiquitination. See, for instance, *PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES*, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993) and Wold, F., *Posttranslational Protein Modifications: Perspectives and Prospects*, pgs. 1-12 in *POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS*, B. C. Johnson, Ed.,

Academic Press, New York (1983); Seifter et al., *Meth. Enzymol.* 182:626-646 (1990) and Rattan et al., *Protein Synthesis: Posttranslational Modifications and Aging*, Ann. N.Y. Acad. Sci. 663: 48-62 (1992). Polypeptides may be branched or cyclic, with or without branching. Cyclic, branched and branched circular polypeptides may result from post-translational natural processes and may be made by entirely synthetic methods, as well.

"Variant(s)" as the term is used herein, is a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. "A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques, by direct synthesis, and by other recombinant methods known to skilled artisans.

EXAMPLES

The examples below are carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. The examples are illustrative, but do not limit the invention.

Example 1 Strain selection, Library Production and Sequencing

The polynucleotide having a DNA sequence given in Table 1 [SEQ ID NO:1] was obtained from a library of clones of chromosomal DNA of *Staphylococcus aureus* in *E. coli*. The sequencing data from two or more clones containing overlapping *Staphylococcus aureus* DNAs was used to construct the contiguous DNA sequence in SEQ ID NO:1. Libraries may be prepared by routine methods, for example:

Methods 1 and 2 below.

Total cellular DNA is isolated from *Staphylococcus aureus* WCUH 29 according to standard procedures and size-fractionated by either of two methods.

Method 1

5 Total cellular DNA is mechanically sheared by passage through a needle in order to size-fractionate according to standard procedures. DNA fragments of up to 11kbp in size are rendered blunt by treatment with exonuclease and DNA polymerase, and EcoRI linkers added. Fragments are ligated into the vector Lambda ZapII that has been cut with EcoRI, the library packaged by standard procedures and *E.coli* infected with the packaged library. The library is amplified by standard procedures.

10 Method 2

Total cellular DNA is partially hydrolyzed with a one or a combination of restriction enzymes appropriate to generate a series of fragments for cloning into library vectors (e.g., RsaI, PstI, AluI, Bsh1235I), and such fragments are size-fractionated according to standard procedures. EcoRI linkers are ligated to the DNA and the fragments then ligated into the
15 vector Lambda ZapII that have been cut with EcoRI, the library packaged by standard procedures, and *E.coli* infected with the packaged library. The library is amplified by standard procedures.

Example 2 Novel ABC transporter Characterization

A family of related ATP-binding proteins carry out distinct biological functions in both eukaryotes and prokaryotes. The majority of proteins in this family transport small molecules across cytoplasmic membranes. All of these proteins share a consensus ATP binding motif of [AG]-x(4)-G-K-[ST]. This family of proteins is known as ABC transporters.

What is claimed is:

1. An isolated polynucleotide comprising a polynucleotide having at least a 70% identity to a polynucleotide encoding a polypeptide comprising the amino acid sequence of SEQ ID NO:2.
2. An isolated polynucleotide comprising a polynucleotide having at least a 70% identity to a polynucleotide encoding the same mature polypeptide expressed by the Novel ABC transporter gene contained in the *Staphylococcus aureus* of the deposited strain.
3. An isolated polynucleotide comprising a polynucleotide encoding a polypeptide comprising an amino acid sequence which is at least 70% identical to the amino acid sequence of SEQ ID NO:2.
4. An isolated polynucleotide that is complementary to the polynucleotide of claim 1
5. The polynucleotide of Claim 1 wherein the polynucleotide is DNA or RNA
6. The polynucleotide of Claim 1 comprising the nucleic acid sequence set forth in SEQ ID NO:1.
7. The polynucleotide of Claim 1 comprising nucleotide 2655 to the stop codon which begins at nucleotide number 3354 set forth in SEQ ID NO:1.
8. The polynucleotide of Claim 1 which encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:2.
9. A vector comprising the polynucleotide of Claim 1.
10. A host cell comprising the vector of Claim 9.
11. A process for producing a polypeptide comprising: expressing from the host cell of Claim 10 a polypeptide encoded by said DNA.
12. A process for producing a Novel ABC transporter polypeptide or fragment comprising culturing a host of claim 10 under conditions sufficient for the production of said polypeptide or fragment.
13. A polypeptide comprising an amino acid sequence which is at least 70% identical to the amino acid sequence of SEQ ID NO:2.
14. A polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:2.
15. An antibody against the polypeptide of claim 14.
16. An antagonist which inhibits the activity or expression of the polypeptide of claim 14.

17. A method for the treatment of an individual in need of Novel ABC transporter polypeptide comprising: administering to the individual a therapeutically effective amount of the polypeptide of claim 14.

18. A method for the treatment of an individual having need to inhibit Novel ABC transporter polypeptide comprising: administering to the individual a therapeutically effective amount of the antagonist of Claim 14.

19. A process for diagnosing a disease related to expression or activity of the polypeptide of claim 14 in an individual comprising:

- (a) determining a nucleic acid sequence encoding said polypeptide, and/or
- (b) analyzing for the presence or amount of said polypeptide in a sample derived from the individual.

20. A method for identifying compounds which interact with and inhibit or activate an activity of the polypeptide of claim 14 comprising:

contacting a composition comprising the polypeptide with the compound to be screened under conditions to permit interaction between the compound and the polypeptide to assess the interaction of a compound, such interaction being associated with a second component capable of providing a detectable signal in response to the interaction of the polypeptide with the compound;

and determining whether the compound interacts with and activates or inhibits an activity of the polypeptide by detecting the presence or absence of a signal generated from the interaction of the compound with the polypeptide.

21. A method for inducing an immunological response in a mammal which comprises inoculating the mammal with Novel ABC transporter polypeptide of claim 14, or a fragment or variant thereof, adequate to produce antibody and/or T cell immune response to protect said animal from disease.

22. A method of inducing immunological response in a mammal which comprises delivering a nucleic acid vector to direct expression of Novel ABC transporter polypeptide of claim 14, or fragment or a variant thereof, for expressing said Novel ABC transporter polypeptide, or a fragment or a variant thereof *in vivo* in order to induce an immunological response to produce antibody and/ or T cell immune response to protect said animal from disease.

SEQUENCE LISTING

<110> SmithKline Beecham Corporation

<120> Novel ABC transporter

<130> GM10125

<150> 08/993,825

<151> 1997-12-18,

<160> 4

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 4279

<212> DNA

<213> Staphylococcus aureus

<400> 1

```

catttatata ttgttgtctt aatgttttat cagaagtttg ttgtaactcg cttaataaat
60
gatgaatatg tgtaaatggt gttttgattt catgagatac attttgtaca aaatgctgcc
120
tcatttgatc aacttgtccc aatgactctc tcattcttatt aaagtgatac tgtaatgttc
180
caatttcac tcgtgctggt tgtttgatag gtgtttcaaa atcaccatca attaagcgtt
240
cggtcgctag cttaacttt tttactggac gaatgattga ataggttgat gcgataacta
300
atgaaatcga gataaataat aacaacatta acaagacggc taagaatggt ctaaactcgc
360
taaagtgttc tccaatatct ggacgcatga atactgctat tgaaccgtct tttgtcttaa
420
agttaatacc aacagtatta tctgttacat tgtcgaaaaa tccagttacg aataatgcga
480

```

atgtttatct ttaataccat ggtaatcttt attattttaat acgtttgtaa tagcattttg
540
tgaaaaagta tcttctctaa atggctcccc aaaaatgtct tatgaccctt ttgatcaacg
600
gtcataatth ggtaattcat ttgcctctaa tgtttgaaat attgttgaat gtgcgttggt
660
ttaactgatt gttcatattg tcttgcttct ttaagtgtct tcatgatttt cgcgtcatta
720
gatgctttta aattataatg atagtaaaca tttgttaata caaaacttat taatgcacta
780
aataaaataa ccgtaatgga ataaatcgca attctagcat agagtgtttt aaacatgatt
840
ctccacctta tagccttgct ctcttactgt ttcaattgta agtgtggcat ttaatttttt
900
taatctttgg cgtagtcgct taatatgaac gtcaactggt cgttcactct cttcataatc
960
atagcccaa attttttcta ttatttggtc acgagtaaag atttgcttag gacgtgctgc
1020
aagcataaat aataattgaa attccttggt cggtaacgct atcgttttatt agatacttgg
1080
agttccaaat aggattgggt tagcgtaaag ttgccaatag tcatttctga atttgaattg
1140
atattatata gacgtaatac cgcacgaatt ctaaaaataa gttccttaac ctcaaagggt
1200
ttggttacat aatcgtcagt accacttata aacgcacgct ctttgctact aagtgcattc
1260
cgcgtgtta acataataac tggatatca taatcatttt ttaatgtatt acataattga
1320
aagccgtcca taccatccat cataatatct accactgcaa tatcgacacg ctgtttttct
1380
aaaagtthta atgcagcttc accacttggt tgtgtgtatg catcaatgtg ctctgtttgt
1440
aatggctag ctatataatt taggattcga ggatcgatc cgacaacaag acattgcacc
1500
atagctataa actcccttat ctttttcatt tattatacat gtaaaatatt tttgcgtaaa
1560
aaaacaattg ttcattattg gttcatattt caaccttata ctgacgctaa agaagaaata
1620
gggagaagtg aatcgatatg aaattagcga taaaagagat tatgttttac aaatttcgtt
1680

atattttaat cacattaatc attcttttat taagtattat ggggttattt attagtgggt
1740
taactcaagg gcttggtagg gagaatattt cgttatttga acactttgat aatgaataat
1800
atgttggtca aaaaatgaaa gagccgcaaa ttgagaaatc gcaacttagt gatacgcaac
1860
agaatcaatt aaaaaagtaa ttcatacaaga accatataaa atgaatattc aaacattgaa
1920
attatctaataaagaacaag atgtgattac atgaatgacg ttaaacaaca acgcctacaa
1980
cttaaaaaag gtgaatatcc taaaaatgct catgaggctg ctattaatga taagttagct
2040
gcagacaaca ttagagtcgg ggatagatta cattttaaaa ataattcaac tagttataga
2100
gtttctggta ttttaaacga cacaatgtat gcgcatagtt ccattgtgct attgaacgat
2160
aacggattta atgcattgaa taaggttaat acggcatttt atccagtga aaatttaaca
2220
caacaacaac gtgatgagct taataaaata aatgacgttc aagttgtgag tgaaaaagat
2280
ttaacaggta atattgagag ttatcaagca gagcaagcac cgttaaatat gatgattgtt
2340
agtttgtttg ctattacagc aatcgttcta agtgcatttt tctatgttat gacgattcaa
2400
aaaatatcac aaattggcat tttgaaagca attggtatta agacaagaca tttattgagt
2460
gcgttagttt taaaaatttt aacactaaca ataattggg taggtattgc tgtgatcatc
2520
atagtaggac tatcatttat gatgccggtg acgatgcctt tttacttaac aacgcaaaat
2580
attttattaa tgggtgggat atttatatta gtagcgattt taggtgcctc actatcattt
2640
atcaaattat ttaaagtgga tcctatcgaa gcaattggag gtgcagaata atggcattag
2700
tcgttaaaga tatcgtcaaa aatttcggag aaggtttgtc tgaaacaaaa gttttaaaag
2760
gtattaattt tgaagtggaa caaggggaat ttgtcatttt aaatggtgcc tctggttctg
2820
ggaaaacaac attgctaacg atattaggcg gattgttaag tcaaacgagt ggtacagtgc
2880

tttacaatga tgcaccattg tttgataaac agcatcgtcc tagtgatttg cgattggaag
2940
atattgggtt ttttttcaa tcttcacatt tagttcctta tttaaaagtg atagagcaat
3000
tgacactcgt aggccaagaa gcgggaatga ccaaacaaca aagttcaaca agagcaatac
3060
aacttttgaa aaatattggg ttagaagatc gcttgaatgt atatccgcat cagttatctg
3120
gcggtgaaaa gcaacgtgtt gcgattatga gagcatttat gaataatccg aaaatcattt
3180
tagcagatga gcccacagca agtttagatg ccgatagagc aacaaaagtt gttgagatga
3240
tacgtcaaca aattaaagaa caacaaatga ttggtattat gattacacac gatcgaagat
3300
tatttgaata tgcagatcga gtgattgaat tagaagatgg caaaataact gattagtggc
3360
ttgtaaagac gctaaatgtt aatgatttaa gacatagtag tataaaagtt agataacaga
3420
atacgatttg gggtttacaaa aaagcgcgtc actattatat aaaatgatga cgtgctttag
3480
tttatttatt ttcaatataa caattgtcta atcaaggatt aacgaatatt taaagatagt
3540
ttgacgcaat attagaaaca acctataata atagtttggt tgtggattaa ctattataaa
3600
taaaagcggc gtaaagacat ataaaccaac tacttgaaca atataacgtt aataacaatc
3660
tatactgata cattacgcct agataatctt taatgagcac atgtaagaaa aagtgatatg
3720
gtgtatgact tccgacacca tcgatagata aacctaatTT ttgggctagt cgtaaggcgc
3780
gcaatacatg aaactgactt gttacacaaa caattttaac tgcttcatga tacaaattgt
3840
tgatgatttg tttagaatat aaaaagtttg tgtatgtatt tatagagtga gattccatta
3900
gtatatctgt tttatcaaca ccatgtgcaa tcaaataacg ttgcatagct aaagcttcag
3960
aaattgggtc gtctggctct tgtccgccag atacaataat ctttggtgct gatgcttggt
4020
gttgatagat atcaagtga cggctaaac gcgctgcaag cattggtgtg acaaattcgg
4080

taaaaataacc agcacctaac acaattatga tatcaacttc tttgttgat gatctatgtc
 4140
 tatatgatac tgtccaaacg agataacaaa taaaggtag taacagggaa agacataata
 4200
 tagctaacca catagacaaa cctttcacia taggtgactg aatcgactt ataaatagaa
 4260
 gtgctgatgt gtagagtac
 4279

<210> 2

<211> 233

<212> PRT

<213> Staphylococcus aureus

<400> 2

Ser	Gly	Ser	Tyr	Arg	Ser	Asn	Trp	Arg	Cys	Arg	Ile	Met	Ala	Leu	Val
1				5					10					15	
Val	Lys	Asp	Ile	Val	Lys	Asn	Phe	Gly	Glu	Gly	Leu	Ser	Glu	Thr	Lys
			20					25					30		
Val	Leu	Lys	Gly	Ile	Asn	Phe	Glu	Val	Glu	Gln	Gly	Glu	Phe	Val	Ile
		35				40						45			
Leu	Asn	Gly	Ala	Ser	Gly	Ser	Gly	Lys	Thr	Thr	Leu	Leu	Thr	Ile	Leu
	50					55					60				
Gly	Gly	Leu	Leu	Ser	Gln	Thr	Ser	Gly	Thr	Val	Leu	Tyr	Asn	Asp	Ala
65					70				75					80	
Pro	Leu	Phe	Asp	Lys	Gln	His	Arg	Pro	Ser	Asp	Leu	Arg	Leu	Glu	Asp
			85					90					95		
Ile	Gly	Phe	Ile	Phe	Gln	Ser	Ser	His	Leu	Val	Pro	Tyr	Leu	Lys	Val
		100						105					110		
Ile	Glu	Gln	Leu	Thr	Leu	Val	Gly	Gln	Glu	Ala	Gly	Met	Thr	Lys	Gln
	115						120					125			
Gln	Ser	Ser	Thr	Arg	Ala	Ile	Gln	Leu	Leu	Lys	Asn	Ile	Gly	Leu	Glu
	130					135					140				
Asp	Arg	Leu	Asn	Val	Tyr	Pro	His	Gln	Leu	Ser	Gly	Gly	Glu	Lys	Gln
145				150					155					160	
Arg	Val	Ala	Ile	Met	Arg	Ala	Phe	Met	Asn	Asn	Pro	Lys	Ile	Ile	Leu
			165					170					175		
Ala	Asp	Glu	Pro	Thr	Ala	Ser	Leu	Asp	Ala	Asp	Arg	Ala	Thr	Lys	Val
	180						185					190			
Val	Glu	Met	Ile	Arg	Gln	Gln	Ile	Lys	Glu	Gln	Gln	Met	Ile	Gly	Ile
	195						200					205			

Met Ile Thr His Asp Arg Arg Leu Phe Glu Tyr Ala Asp Arg Val Ile
210 215 220
Glu Leu Glu Asp Gly Lys Ile Thr Asp
225 230

<210> 3

<211> 20

<212> DNA

<213> Staphylococcus aureus

<400> 3

tgcctctggt tctgggaaaa

20

<210> 4

<211> 33

<212> DNA

<213> Staphylococcus aureus

<400> 4

actaaatgtg aagattgaaa aataaaacca ata

33

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/26037

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C07K 14/31; C12N 01/21, 15/31, 15/63, 15/74

US CL : 530/350, 825; 536/23.1, 23.7; 435/69.1, 252.3, 320.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 530/350, 825; 536/23.1, 23.7; 435/69.1, 252.3, 320.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, MEDLINE, BIOSIS, MasPar

search terms: ABC transporter#, Staphylococc## aureus

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- A	MALDENER et al. Characterization of devA, a Gene Required for the Maturation of Proheterocysts in the Cyanobacterium Anabaena sp. Strain PCC 7120. Journal of Bacteriology. December 1994, Vol. 176, No. 24, pages 7543-7549, especially pages 7545-7547.	4 and 12 ----- 1-3, 5-11 and 13-14
X --- A	WO 97/23620 A1 (SMITHKLINE BEECHAM P.L.C.) 03 July 1997, pages 12, 18-20, 23-31 and Figure 1.	2, 4 and 12 ----- 1, 3, 5-11 and 13-14



Further documents are listed in the continuation of Box C.



See patent family annex.

* "A"	Special categories of cited documents: document defining the general state of the art which is not considered to be of particular relevance	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
E	earlier document published on or after the international filing date	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
L	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
O	document referring to an oral disclosure, use, exhibition or other means	
P	document published prior to the international filing date but later than the priority date claimed	*A* document member of the same patent family

Date of the actual completion of the international search

22 MARCH 1999

Date of mailing of the international search report

19 APR 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

ROBERT C. HAYES, PH.D.

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/26037

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-14

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/26037

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-14, drawn to a polynucleotide, ABC transporter polypeptide, vector, host cells and method for producing the polypeptide.

Group II, claim(s) 15, drawn to an antibody.

Group III, claim(s) 16, drawn to an antagonist.

Group IV, claim(s) 17, drawn to a method of treatment comprising administering a polypeptide.

Claim V, claim(s) 18, drawn to a method of treatment comprising administering an antagonist.

Group VI, claim(s) 19(a), drawn to a process for diagnosing a disease state comprising determining a nucleic acid sequence.

Group VII, claim(s) 19(b), drawn to a process for diagnosing a disease state comprising analyzing a polypeptide.

Group VIII, claim(s) 20, drawn to a method for identifying compounds which interact with a polypeptide.

Group IX, claim(s) 21, drawn to a method for inducing an immunological response which comprises inoculation with a polypeptide.

Group X, claim(s) 22, drawn to a method for inducing an immunological response which comprises delivering a nucleic acid vector.

The inventions listed as Groups I-X do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

Group I is directed to a polynucleotide, ABC transporter polypeptide, vector, host cells and method for producing the polypeptide, which is the first product and first method of using the product. The special technical feature is the polynucleotide encoding the ABC transporter. Groups I-III are drawn to structurally different products, which do not require each other for their practice and do not share the same or a corresponding technical feature. The Group IV-X inventions are drawn to methods having different goals, method steps and starting materials, which do not require each other for their practice and do not share the same or a corresponding technical feature. Note that PCT Rule 13 does not provide for multiple products or methods within a single application. Since the special technical feature of the Group I invention is not present in the Group II-X claims, and the special technical features of the Group II-X inventions are not present in the Group I claims, unity of invention is lacking.